



SIGMOIDOSCOPY OR SHORT COLONOSCOPY

YOUR EXAM

Your doctor has prescribed a sigmoidoscopy exam which is a short colonoscopy exam. In this procedure a flexible tube is introduced through the anus. The tube has a camera and light source that allows the physician to examine the large intestine and detect the presence of lesions. The most common type of lesions is called polyps^[1].

During the procedure, your doctor may take a biopsy^[2] or remove the polyps.

[1] Polyps are benign tumours which may measure from several millimeters to several centimeters. They should be removed because the lesions can become cancerous.

[2] A medical test involving the removal of tissues in order to analyze and to determine their nature.

WHAT TO DO BEFORE YOUR EXAM

It is important to have an enema and to follow the recommended guidelines, because the quality and safety of your exam depend on it.

- If you are taking iron, Coumadin (Warfarin), Plavix, Aspirin, or Heparin, notify the clinical staff when scheduling your appointment.
- Bring an up-to-date list of all medications you are taking and be sure to mention any known allergies.
- You may eat and drink normally.
- Do not wear jewellery or nail polish the day of the exam.
- Arrive 30 minutes before the scheduled time of your exam.

IMPORTANT

You will not be able to drive after the exam, so we ask that you be accompanied by a friend or family member.

DURING THE EXAM

- The exam may last from 10 to 20 minutes.
- You will be moved to an examination stretcher.
- You will lie on your left side.
- The nurse will give you a mild sedative to keep you comfortable and help you relax.
- When you will be ready, the doctor will introduce the tube into your anus. It is important to mention if you feel any pain.

AFTER THE EXAM

- You will be kept in an observation area for about 30 minutes.
- The staff will regularly check your blood pressure and other vital signs such as your heart rate.
- It is normal to feel bloated or have gas.
- It is normal to feel tired and nauseous.

AT HOME

The overall risk of complications with this exam is very low:

- bleeding;
- gastrointestinal perforation.

If you have severe stomach pains, chills, or bleeding:



Consult with your family doctor or go to a walk-in health clinic.



If the clinic is closed:
call Info-Santé 8-1-1.



Go to the nearest hospital emergency department.

Authors

Frédéric Grondin, nurse, M.Sc.; Dr Charles Ménard, gastroenterologist

Revision and layout

Service des communications

Direction des ressources humaines, des communications et des affaires juridiques

© Centre intégré universitaire de santé et de services sociaux de l'Estrie – Centre hospitalier universitaire de Sherbrooke, 2017

santeestrie.qc.ca

July 2017 – 1-6-70928

The information provided in this document is made possible thanks to financial support from the Fonds Brigitte-Perreault for patient safety and quality of care.

If you must cancel your exam, please notify the staff at 819 346-1110, ext 24462.