



PRECAUTION | ANALYSIS OF URINARY PROSTAGLANDIN D2 OR E2

This test is used to screen for certain health problems or diseases and helps to verify the level of prostaglandin in your urine.

PREPARATION

The presence of certain substances in your urine can cause falsely elevated results. Therefore, **two days (48 hours) before and on the day of the urine collection, do not take the following medications: Aspirin®, indometacin, anti-inflammatory drugs.** However, you must first speak to your prescriber before you stop taking any prescribed medications.

DO YOU HAVE ANY QUESTIONS?



santeestrie.qc.ca/sampling



Info-Santé 811
OR
Speak to your prescriber.

Authors

Laboratory medicine team

Approval

Dr. François Goyer, biochemist

Revision and layout

Service des communications

Direction des ressources humaines, des communications et des affaires juridiques

© Centre intégré universitaire de santé et de services sociaux
de l'Estrie – Centre hospitalier universitaire de Sherbrooke 2020

Document management code | INS-CP-3097-FR

santeestrie.qc.ca

June 2023 – 1-6-72034 (French version) | 1-6-72035 (English version)

**Centre intégré
universitaire de santé
et de services sociaux
de l'Estrie – Centre
hospitalier universitaire
de Sherbrooke**

Québec 