



Avec vous, pour la Vie

URODYNAMIC ASSESSMENT

A urodynamic assessment will allow your physician to evaluate how your bladder and sphincter function, and understand your problem with storage or emptying of urine.

WHO SHOULD BE TESTED?

Persons:

- Suffering from urinary incontinence, interstitial cystitis, urinary infection, benign hypertrophy of the prostate, or other disorders of the urinary system.
- Receiving intravesical treatment (BCG or other).
- Who have difficulty urinating or whose bladder does not empty completely.

WHAT SHOULD YOU DO BEFORE THE TEST?

- No preparation is required in the days prior to the test.
- You do not need to change or stop taking your medication.
- You can have breakfast or lunch before the test.
- This test carries little risk and is not painful.
- It is normal to feel some apprehension before the test. This uneasiness should lessen during the test.

DAY OF THE TEST


- Drink two glasses of water one hour prior to the test and do not urinate before you have the test.
- If you wear a urinary catheter, you need to clamp it one hour before the test to stop the flow of urine.
- You must have a full bladder when you arrive for your first test.
- Wear a two-piece outfit because you'll need to remove the clothing from your lower body.
- Go to the letter G reception desk at the outpatient clinics at the CHUS – Fleurimont Hospital (3001, 12th Avenue N., Sherbrooke) to indicate your arrival.

You will then be asked to go to the letter H waiting room where you will be greeted by the urodynamic room staff.

- Bring the most up-to-date list of medications you take.
- Notify the urodynamic room staff if you recently had a urinary infection, whether it was treated or untreated, or if you believe you currently have one.
- The test takes about 60 minutes.

DURING THE TEST

The tests described below will help your physician better understand how your bladder functions.



- 1. **Uroflowmetry: You must arrive with a full bladder** because we will ask you to urinate into a special toilet linked to a computer that will measure the force and regularity of the flow.
- Cystometrography consists of introducing a narrow catheter through the urethra to fill the bladder with sterile water, making it possible to observe how your bladder reacts when you cough or exert effort.
- Sphincter electromyography (EMG)
 measures the coordinated action of several
 muscles and records sphincter muscle activity
 during filling and emptying of the bladder.
- 4. The **urethral profile** measures the pressure variations along the entire length of the urethra.

AFTER THE TEST

- Drink lots of water to reduce the irritation caused by the catheter in the bladder.
- The urologist or your family physician will provide the results during a medical appointment.

ONCE YOU'RE HOME

- If you experience severe abdominal pain;
- If you have a fever (more than 38.5°C), with or without chills;
- Or in the event of major bleeding or persistent burning when urinating;



Contact your family physician and immediately tell him or her that you had a urodynamic assessment.



If you cannot reach your physician, call Info-Santé 811.



Or go to the emergency department of the hospital nearest you.

Source

- Josée Parenteau, Clinical Nurse | General and Emergency Medicine Client Program
- Renée Létourneau, Clinical Nurse in Clinical Development | Surgical Client Care Program

Review

- Gaëlle Simon, AIC | Outpatient Clinics | General and Emergency Medicine Client Care Program
- DCAP

Layout: DCAP

Illustration: Canadian Urological Association

© Centre hospitalier universitaire de Sherbrooke (CHUS) chus.gc.ca

September 2013 - 1-6-71723

REMINDER

Date of your test:

Should you need to cancel your test, please notify the administrative officer at 819-564-5210 or 1-866-638-2601.

CHUS – Fleurimont Hospital

Urodynamic Room

Room 26H4, Letter G, Outpatient Clinics 3001 - 12th Avenue N., 2nd floor Sherbrooke, Quebec J1H 5N4