

# **CIRCUMCISION**

#### **ABOUT YOUR SURGERY**

Posthectomy, commonly known as circumcision, is a procedure involving the removal of the foreskin from the tip of the penis. Two types of circumcision are available depending on your medical condition: full and partial.

### **BEFORE YOUR SURGERY**

- Inform the physician or nurse if you are taking blood thinners (anticoagulants) or anti-inflammatory drugs.
- The evening before the procedure, shave your scrotum and pubic hair.

## **PROCEDURE**

- The surgery is usually performed under local anesthesia among adults.
- The procedure lasts approximately 20 to 30 minutes.
- Once the foreskin is excised, dissolvable stitches are applied to close the wound.
- A Bactigras®-type dressing will be applied. The dressing contains petroleum jelly (Vaseline®) and other agents to prevent infection.
- You will be kept in the observation room for 15 minutes.

# **AFTER THE SURGERY**

- You can walk, and climb up and down stairs as soon as the procedure has ended.
- You can shower 24 hours later.
- Do not engage in physical activity for 1 week.
- Avoid baths, spas, and swimming pools for 1 week.
- Nocturnal erections will be very painful, but once you are awake and the erection has ended, so will the pain.
- Sexual intercourse is permitted after 4 to 6 weeks.
- You can go back to work when recommended by your urologist.



# **POTENTIAL SIDE EFFECTS**

- Significant pain for the first 2 days after the procedure. It is recommended to regularly take 2 acetaminophen (Tylenol®) tablets, 4 times daily, then as needed. If the pain persists, you can take prescribed narcotics.
- Bruising and edema (swelling) should go away after 2 months.
- Mild oozing of pus or blood around the wound.



An appointment will be scheduled with your doctor in the months after your surgery.

### WHEN TO CONSULT

If you have any concerns regarding your health status or if you notice any of the following symptoms:

- Purulent discharge around the wound
- Fever/chills
- Increasing pain after 1 week
- Abnormal bleeding

Call Info-Santé 8-1-1 or the urology reception desk at 819 346-1110, Ext. 74595.

#### Authors

Gabrielle Lalumière-Houde nurse clinician, Dr Robert Sabbagh, urologist, and Dr Claudio Jeldres, urologist.

#### Revision and layout

Service des communications

Direction des ressources humaines, des communications et des affaires juridiques

© Centre intégré universitaire de santé et de services sociaux de l'Estrie – Centre hospitalier universitaire de Sherbrooke 2022

santeestrie.qc.ca

October 2022 - 4-6-11517 (French version) | 4-6-11518 (English version)