

HYALURONIDASE (MEDICATION) - CHILD

Your child has been prescribed this medication to reduce potential damage caused by a chemotherapy leak out of his vein and beneath his skin. This is known as an extravasation.

POTENTIAL SIDE EFFECTS

- Swelling at the injection site
- Allergic reaction (rash, itching, swelling around the eyes)
- Local irritation
- Infection
- Bleeding
- Bruising

Nurse's signature:

Date: _____

Telephone number and extension:

WHEN TO CONSULT

Contact the nurse at the pediatric oncology centre **as soon as possible** if you notice:

- Onset of or increasing pain
- Onset of or increasing redness
- Onset of itching
- Presence of a skin lesion (blister)
- Skin breakdown
- Presence of discharge
- Swelling
- Difficulty moving the hand, wrist, elbow or shoulder of the arm where the extravasation occurred

Notify the nurse at the pediatric oncology centre and go to the emergency department if you notice:

- Swelling around the eyes.
 - Fever of 38.3 °C (101 °F) once.
- OR
- Fever of 38 °C (100.4 °F) twice within a one-hour interval.

Editing

Direction des soins infirmiers

Adapted from *Guide de prise en charge de l'extravasation des agents antinéoplasiques*. INESSS, April 2019.

Revision and layout

Service des communications

Direction des ressources humaines, des communications et des affaires juridiques

© Centre intégré universitaire de santé et de services sociaux de l'Estrie – Centre hospitalier universitaire de Sherbrooke 2021

santeestrie.qc.ca

February 2022 – 4-6-11200 (French version) | 4-6-11201 (English version)

Centre intégré
universitaire de santé
et de services sociaux
de l'Estrie – Centre
hospitalier universitaire
de Sherbrooke

Québec

