



## END OF RADIATION THERAPY TREATMENTS | PROSTATE

You have just had your last radiation therapy treatment. The side effects (redness or bump on your tattoos, redness at the area treated, diarrhea or frequent bowel movements, frequent urge to urinate, burning sensation when urinating, fatigue) may persist or even increase during the two weeks after your treatments are over. These effects will gradually disappear.

### RECOMMENDATIONS TO FOLLOW AT THE END OF TREATMENTS

In the coming weeks, it will be important to follow these recommendations to minimize the complications that may occur.

#### Caring for the skin at the area treated

- Be gentle with your skin.

#### Diarrhea or frequent bowel movements

Even though your treatments have ended, you may experience diarrhea or frequent small bowel movements. Should this be the case, return to the low-fibre diet recommended during your treatments. This situation should resolve itself in a few weeks. Then, return to a balanced diet gradually.

**Return to daily routine activities:** return to your regular activities progressively.

#### In the event of complications or a worsening of the side effects associated with radiation therapy



You can reach the Radiation Oncology Department at 819-346-1110, ext. 14602, Monday to Friday, 8 a.m. to 4 p.m.



If the Department is closed, call Info-Santé at 8-1-1.



Consult your family doctor or go to emergency at the nearest hospital.

#### Author

Service de radio-oncologie de l'Hôpital Fleurimont

#### Revision and layout

Direction des ressources humaines, des communications et des affaires juridiques

© CIUSSS de l'Estrie – CHUS

[chus.qc.ca](http://chus.qc.ca)

August 2015 – 1-6-71826